

On the Facets of Mixed Integer Programs with Two Integer Variables and Two Constraints

G erard Cornu ejols *

Tepper School of Business, Carnegie Mellon University, Pittsburgh, PA 15213
and LIF, Facult e des Sciences de Luminy, Universit e de Marseille, France
gc0v@andrew.cmu.edu

and

Fran ois Margot †

Tepper School of Business, Carnegie Mellon University, Pittsburgh, PA 15213
fmargot@andrew.cmu.edu

August 2007; revised March 2008

Abstract

In this paper we consider an infinite relaxation of the mixed integer linear program with two integer variables, nonnegative continuous variables and two equality constraints, and we give a complete characterization of its facets. We also derive an analogous characterization of the facets of the underlying finite integer program.

1 Introduction

We consider the mixed 2-integer-variable linear program with two constraints

$$\begin{aligned}x &= f + \sum_{j=1}^k r^j s_j \\x &\in \mathbb{Z}^2 \\s &\in \mathbb{R}_+^k\end{aligned}\tag{1}$$

where $f \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$, $k \geq 1$, and $r^j \in \mathbb{Q}^2 \setminus \{0\}$. Let $R_f(r^1, \dots, r^k)$ be the convex hull of all vectors $s \in \mathbb{R}_+^k$ such that $f + \sum_{j=1}^k r^j s_j$ is integral. $R_f(r^1, \dots, r^k)$ is a polyhedron (We refer the reader to [17] for standard definitions). Model (1) was considered by Andersen, Louveaux, Weismantel and Wolsey [1]. They showed that the nontrivial facets of $R_f(r^1, \dots, r^k)$ are necessarily defined by split inequalities or intersection cuts (Balas [2]) arising from triangles or quadrilaterals in \mathbb{R}^2 . A goal of this paper is to give a converse to the result in [1]: which splits, triangles and quadrilaterals actually define facets of $R_f(r^1, \dots, r^k)$? We present our analysis in the more general context of the Gomory-Johnson infinite group relaxation.

*Supported by NSF grant CMMI0653419, ONR grant N00014-97-1-0196 and ANR grant BLAN06-1-138894.

†Supported by ONR grant N00014-97-1-0196.

Gomory [9], Gomory and Johnson [12] and Johnson [13] suggested relaxing the k -dimensional space of variables $s = (s_1, \dots, s_k)$ to an infinite-dimensional space, where the variables s_r are defined for any $r \in \mathbb{Q}^2$. We get the *infinite program with two integer variables and two constraints*

$$\begin{aligned} x &= f + \sum r s_r \\ x &\in \mathbb{Z}^2 \\ s &\geq 0 \quad \text{with finite support.} \end{aligned} \tag{2}$$

The vector $s = (s_r)_{r \in \mathbb{Q}^2}$ is said to have *finite support* if $s_r \neq 0$ for a finite number of $r \in \mathbb{Q}^2$. Let R_f be the convex hull of all vectors $s \geq 0$ with finite support such that $f + \sum r s_r$ is integral. Note that the polyhedron $R_f(r^1, \dots, r^k)$ is obtained from R_f by setting $s_r = 0$ for all $r \in \mathbb{Q}^2 \setminus \{r^1, \dots, r^k\}$. Our motivation for working with R_f is that it only has one parameter, namely f , and therefore the results are cleaner than with $R_f(r^1, \dots, r^k)$. A drawback of R_f is that it is not a closed set. For example, the sequence s^k for $k = 1, 2, \dots$ defined by

$$s_r^k = \begin{cases} \frac{1}{k} & \text{if } r = -kf \\ 0 & \text{otherwise} \end{cases}$$

is in R_f and converges to 0, but $0 \notin R_f$ since $f \in \mathbb{Q}^2 \setminus \mathbb{Z}^2$. Throughout the paper, we relate results obtained for the convex set R_f to those obtained for the polyhedron $R_f(r^1, \dots, r^k)$.

1.1 Minimal valid inequalities

We say that an inequality $\alpha s \geq \beta$ is *valid* for R_f (resp. $R_f(r^1, \dots, r^k)$) if it is satisfied by all vectors in R_f (resp. $R_f(r^1, \dots, r^k)$). Inequalities $s_i \geq 0$ are called *trivial* valid inequalities. In this paper, we discuss only nontrivial valid inequalities. The solution $s = 0$ is not feasible for R_f . Any valid inequality for R_f that cuts off the vector $s = 0$ is of the form

$$\sum \psi(r) s_r \geq 1, \tag{3}$$

where $\psi : \mathbb{Q}^2 \rightarrow \mathbb{R} \cup \{+\infty\}$ and, as above, we only consider vectors s with finite support. To avoid ambiguity, the product $+\infty \cdot 0$ is defined to be 0.

Any valid inequality for R_f yields a valid inequality for $R_f(r^1, \dots, r^k)$ by simply restricting it to the space r^1, \dots, r^k . Furthermore, a full description of the polyhedron $R_f(r^1, \dots, r^k)$ is obtained from the set of valid inequalities for R_f by adding the constraints $s_r = 0$ for $r \neq r^1, \dots, r^k$. Therefore we will assume in the remainder that valid inequalities for $R_f(r^1, \dots, r^k)$ are restrictions of valid inequalities for R_f .

An inequality $\sum \psi(r) s_r \geq 1$ valid for R_f is *minimal* if there is no valid inequality $\sum \psi'(r) s_r \geq 1$ where $\psi' \leq \psi$ and $\psi'(r) < \psi(r)$ for at least one $r \in \mathbb{Q}^2$. Note that when $\psi(r) = +\infty$ we have $\psi'(r) < \psi(r)$ if and only if $\psi'(r)$ is finite. If ψ defines a minimal valid inequality $\sum \psi(r) s_r \geq 1$, we also say that the function ψ is *minimal*. In [4], it was shown that a minimal valid function ψ is nonnegative, positively homogeneous, piecewise linear, and convex. Recall that a function ψ is *positively homogeneous* if $\psi(\lambda r) = \lambda \psi(r)$ for all $\lambda \geq 0$. Since ψ is always nonnegative in this paper, we simply say *homogeneous* to mean positively homogeneous. Define

$$B_\psi := \{x \in \mathbb{Q}^2 : \psi(x - f) \leq 1\}. \tag{4}$$

The convexity of ψ implies that B_ψ is a convex set in \mathbb{Q}^2 . The following result was proved in [4].

Theorem 1.1. *Let $f \in \mathbb{Q}^2$. A minimal valid function ψ for R_f is nonnegative, homogeneous, piecewise linear, and convex. Furthermore, the closure of the set B_ψ in \mathbb{R}^2 is a full-dimensional polyhedron with 2, 3 or 4 edges, it contains no integral point in its interior but each edge contains at least one integral point in its relative interior.*

We will simply say in the *interior* of an edge to mean in its *relative interior*. Next we describe the different sets B_ψ arising in Theorem 1.1.




Figure 1: Representation of B_ψ for nondegenerate cases.

The point f is in B_ψ since $\psi(0) = 0$. The *nondegenerate case* is obtained when f is in the interior of B_ψ (see Figure 1). In this case, an extension of ψ to \mathbb{R}^2 is obtained by defining $\psi(x - f) = 1$ for all points on the boundary of the closure of B_ψ , denoted by $\text{cl}B_\psi$. Indeed, the knowledge of f and of the boundary of $\text{cl}B_\psi$ together with the homogeneity of ψ is enough to compute the value of $\psi(r)$ for any vector $r \in \mathbb{R}^2 \setminus \{0\}$: If there is a positive scalar λ such that the point $f + \lambda r$ is on the boundary of $\text{cl}B_\psi$, we get that $\psi(r) = 1/\lambda$. Otherwise, if there is no such λ , r is an unbounded direction of $\text{cl}B_\psi$ and $\psi(r) = 0$. Note that this extension of ψ to \mathbb{R}^2 is a continuous function.




Figure 2: Level curves of $\psi(r)$ for nondegenerate cases.

We use the graphic representation of B_ψ to describe ψ when possible. The inequalities corresponding to the three cases of Figure 1 will be called *split*, *triangle* and *quadrilateral* inequalities. They are special cases of the *intersection cuts* of Balas [2]. Split inequalities in higher dimensions were studied by Cook, Kannan and Schrijver [6], who coined the name. These inequalities are equivalent to Gomory's mixed integer inequalities [10] (see [5] for a

proof). In Figure 2, we represent the function ψ in the (r_1, r_2) -space where $r = \begin{pmatrix} r_1 \\ r_2 \end{pmatrix} \in \mathbb{R}^2$. Solid lines give level curves of $\psi(r)$ with values 0 and 1 for the three examples of Figure 1.

The *degenerate case* is obtained when f is a vertex of $\text{cl}B_\psi$ or when f lies on one of its edges (see Figure 3 for three examples). Then ψ is not finite everywhere [4]. An extension of ψ to $\mathbb{R}^2 \cup \{+\infty\}$ is obtained by defining $\psi(x - f) = 1$ for all points on the boundary of $\text{cl}B_\psi$, except for directions $(x - f)$ supporting edges of $\text{cl}B_\psi$ containing f : For any direction $0 \neq r \in \mathbb{R}^2$ such that the half-line $L_r = \{x = f + \lambda r \text{ for } \lambda > 0\}$ is entirely outside $\text{cl}B_\psi$, we have $\psi(r) = +\infty$. For the directions such that the half-line L_r goes through the interior of $\text{cl}B_\psi$, let $f + \lambda r$ be the point where L_r intersects the boundary of $\text{cl}B_\psi$; then we get $\psi(r) = 1/\lambda$. Finally, when L_r supports an edge E of $\text{cl}B_\psi$, assume that $y = f + \lambda r$ is the first integral point encountered on L_r starting from f (such a point exists since E contains an integral point and f is rational, implying that L_r contains infinitely many integral points) and let $x = f + \mu r$ be the first vertex of $\text{cl}B_\psi$ encountered (if any); if y is encountered first, we get $\psi(r) = 1/\lambda$ and if x is encountered first, we get $\psi(r) = 1/\mu$. The resulting extension of ψ to $\mathbb{R}^2 \cup \{+\infty\}$ is not continuous, even in the region where ψ is finite. Dey et al. [7] showed in a more general context that, if $\psi(r) < +\infty$ everywhere, then ψ is continuous, and therefore ψ is nondegenerate.

There are five different degenerate inequalities, depending on the type of set $\text{cl}B_\psi$ and the position of f on its faces: *degenerate split*, *vertex-degenerate triangle*, *edge-degenerate triangle*, *vertex-degenerate quadrilateral* and *edge-degenerate quadrilateral* inequalities. Solid lines in Figure 4 give level curves of $\psi(r)$ with value 1 for the three examples of Figure 3.




Figure 3: Representation of B_ψ for degenerate cases.

A convex set with no integral point in its interior is called *lattice-free*. Maximal lattice-free convex sets in \mathbb{R}^n are polyhedra with at most 2^n facets (Bell [3], Scarf [18] and Lovász [14]). The complete list of all maximal lattice-free convex sets in the plane is known:

Theorem 1.2. (Lovász [14]) *A maximal lattice-free convex set in the (x_1, x_2) -plane \mathbb{R}^2 is one of the following:*

- i) *A line $ax_1 + bx_2 = c$, where a/b is irrational;*
- ii) *A strip $c \leq ax_1 + bx_2 \leq c + 1$ where a and b are coprime integers and c is an integer;*
- iii) *A triangle with a least one integral point in the interior of each of its edges;*




Figure 4: Level curves of $\psi(r)$ for degenerate cases.

- iv) A quadrilateral containing exactly four integral points, with exactly one of them in the interior of each of its edges; Moreover, these four integral points are vertices of a parallelogram of area 1.

The polyhedra referred to in Theorem 1.1 correspond to the last three cases of Theorem 1.2. The first case does not play a role here as we only consider rational vectors f and r in the definition of R_f . Note that if we had defined R_f and $R_f(r^1, \dots, r^k)$ for parameters in \mathbb{R}^2 instead of \mathbb{Q}^2 , the strict inequality $\sum_{r \neq r^0} s_r > 0$ would be valid when r_1^0/r_2^0 is irrational and the line $f + \lambda r^0$ contains no integral point (this inequality corresponds to the first case of Theorem 1.2 and it is not implied by the valid inequalities of Theorem 1.1 since these inequalities all have $\psi(r^0) > 0$). In particular $R_f(r^1, \dots, r^k)$ would not be a polyhedron anymore. Thus the assumption that the vectors f and r are rational is important in the definitions of R_f and $R_f(r^1, \dots, r^k)$.

On the other hand, extending the definition of ψ and B_ψ to \mathbb{R}^2 is useful since the maximal lattice-free triangles and quadrilaterals in cases ii) and iii) of Theorem 1.2 may have irrational corner vertices. In the remainder of the paper, ψ and B_ψ are always assumed to be defined over \mathbb{R}^2 .

1.2 Facets

A valid inequality $\sum \psi(r) s_r \geq 1$ for R_f is *extremal*, or *defines a facet* of R_f if there does not exist two distinct valid inequalities $\sum \psi_j(r) s_r \geq 1$, $j = 1, 2$, such that $\psi = \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$. By extension, we also say that the function ψ itself *defines a facet* of R_f . Note that, although we only use nontrivial inequalities in this definition, including them would give an equivalent definition.

Remark 1.3. *The definition of a facet implies that if ψ defines a facet of R_f then ψ is minimal.*

Remark 1.4. *If a valid function ψ is not facet defining for R_f , then $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_1 and ψ_2 are both minimal valid functions and $\psi \neq \psi_1$ or ψ_2 .*

Gomory [11] recently noted that triangle inequalities define facets of R_f and raised the question of completely describing the facets of R_f . In this paper, we give such a characterization. It is summarized in the next theorem.

Theorem 1.5. *The facets of R_f are all split inequalities, all triangles inequalities, all quadrilateral inequalities that satisfy a certain ratio condition, all degenerate split inequalities, some degenerate triangle inequalities (see Theorems 4.9 and 4.10 for details) but no degenerate quadrilateral inequalities.*

A valid inequality $\sum_{i=1}^k \psi(r^i)s_i \geq 1$ for $R_f(r^1, \dots, r^k)$ defines a facet of $R_f(r^1, \dots, r^k)$ if there does not exist two distinct valid inequalities $\sum_{i=1}^k \psi_j(r^i)s_i \geq 1$, $j = 1, 2$, such that $\psi(r^i) = \frac{1}{2}\psi_1(r^i) + \frac{1}{2}\psi_2(r^i)$ for $i = 1, \dots, k$. This definition of a facet of $R_f(r^1, \dots, r^k)$ is consistent with the usual definition of a facet of a polyhedron only if the polyhedron is full dimensional. The next lemma shows that this is the case.

Lemma 1.6. *If $R_f(r^1, \dots, r^k)$ is non empty, then it is full dimensional.*

Proof. The recession cone of $R_f(r^1, \dots, r^k)$ is \mathbb{R}_+^k . □

In the remainder, we assume that $R_f(r^1, \dots, r^k) \neq \emptyset$. Note that this is easy to check. In particular, this is always the case when the rays r^j for $j = 1, \dots, k$ span \mathbb{R}^2 .

In [1], Andersen, Louveaux, Weismantel and Wolsey study $R_f(r^1, \dots, r^k)$ and they prove that, when nonnegative combinations of r^1, \dots, r^k span \mathbb{R}^2 , all the nontrivial facets of $R_f(r^1, \dots, r^k)$ are split inequalities or are triangle or quadrilateral inequalities where the vertices of B_ψ are on the rays $f + \lambda r^i$, $\lambda > 0$, for $i = 1, \dots, k$. They do not, however, describe precisely which triangles and quadrilaterals generate facets. Some of the polyhedra B_ψ they use are maximal lattice-free but, in order to have vertices of B_ψ on rays $f + \lambda r^i$, $\lambda > 0$, they also use polyhedra B_ψ that are not maximal lattice-free. See Figure 5.




Figure 5: Data, a non maximal lattice-free quadrilateral defining a facet, and a maximal lattice-free triangle defining the same facet.

In addition to Theorem 1.5, the other main result of this paper is a characterization of the facets of $R_f(r^1, \dots, r^k)$.

Theorem 1.7. *The facets of $R_f(r^1, \dots, r^k)$ are*

- i) split inequalities where the infinite direction of B_ψ is r^j for some $j = 1, \dots, k$ and the line $f + \lambda r^j$ contains no integral point; or where B_ψ satisfies a certain ray condition (see Theorem 3.12 for details)*
- ii) triangle inequalities where the triangle B_ψ has its corner points on three half-lines $f + \lambda r^j$ for some $j = 1, \dots, k$ and $\lambda > 0$; or where the triangle B_ψ satisfies a certain ray condition (see Section 3.2.1 for details),*
- iii) quadrilateral inequalities where the corners of B_ψ are on four half-lines $f + \lambda r^j$ for some $j = 1, \dots, k$ and $\lambda > 0$, and B_ψ satisfies a certain ratio condition (see Theorem 3.10 for details).*
- iv) None of the degenerate cases are needed to define the facets of $R_f(r^1, \dots, r^k)$.*

The paper is organized as follows. In Section 2, we give simple technical results that are useful in the remainder of the paper. In Section 3 we prove Theorem 1.7 and in Section 4 we prove Theorem 1.5.

2 Preliminaries

The three lemmas in this section collect simple properties that will be used in the remainder of the paper.

The first lemma is a characterization of maximal lattice-free triangles that was observed by Dey and Wolsey [8].

Lemma 2.1. *If T is a maximal lattice-free triangle in \mathbb{R}^2 , then*

- i) T has exactly three integral points on its boundary, one in the interior of each edge, or*
- ii) T has exactly six integral points on its boundary, the three vertices and one in the middle of each edge, or*
- iii) T has exactly one edge E with at least two integral points in its interior. The vertex opposite edge E is nonintegral.*

Proof. Suppose T has an edge E_1 with at least two integral points y^1, y^2 in its interior, and another edge E_2 with two integral points y^3, y^4 distinct from the point $x^3 = E_1 \cap E_2$. Say x^3, y^1, y^2 appear in this order in E_1 and x^3, y^3, y^4 appear in this order in E_2 . Then the points $y^2 + (y^3 - y^1)$ and $y^4 + (y^1 - y^3)$ are integral points and at least one of them is in the triangle $x^3 y^2 y^4$ and thus in the interior of T , a contradiction. This shows that either iii) holds or every edge of T has exactly one integral point in its interior.

In the latter case, either i) holds or T has an integral corner point x^3 . Let E_1 and E_2 be the edges of T containing x^3 and let y^i be the integral vertex in the interior of E_i for $i = 1, 2$. Let x^i be the symmetric of x^3 relative to y^i . Since x^i is integral and therefore not interior to E_i by our assumption, T is contained in the triangle $x^1 x^2 x^3$. It is easy to verify that the triangle $x^1 x^2 x^3$ only contains six integral points, namely the x^i 's, y^1, y^2 and the middle

y^3 of x^1 and x^2 : Indeed, the only integral points in triangle $x^3y^1y^2$ are its corners and by translation and symmetry the same holds for $x^1y^1y^3$, $y^1y^2y^3$ and $x^2y^2y^3$. By maximality, T is the triangle $x^1x^2x^3$. Therefore ii) holds. \square

Lemma 2.2. *Let $f \in \mathbb{Q}^2$ and let ψ be valid and minimal for R_f . Let $x^1, x^2, x^3 \in \mathbb{R}^2$ be three distinct points such that $x^2 = \mu x^1 + (1 - \mu)x^3$ for some $0 < \mu < 1$.*

- i) *If $\psi(x^1 - f) \leq 1$ and $\psi(x^2 - f) = 1$ then $\psi(x^3 - f) \geq 1$;*
- ii) *If $\psi(x^1 - f) \leq 1$, $\psi(x^2 - f) = 1$, and $\psi(x^3 - f) \leq 1$ then $\psi(x^1 - f) = \psi(x^3 - f) = 1$;*
- iii) *If $\psi(x^1 - f) = \psi(x^2 - f) = \psi(x^3 - f) = 1$ then $\psi(x - f) = 1$ for all x on the line segment $\mu'x^1 + (1 - \mu')x^3$ for $0 \leq \mu' \leq 1$.*

Proof. i) By Theorem 1.1, ψ is convex and thus $1 = \psi(x^2 - f) \leq \mu\psi(x^1 - f) + (1 - \mu)\psi(x^3 - f) \leq \mu + (1 - \mu)\psi(x^3 - f)$.

ii) Using i), we have $\psi(x^3 - f) \geq 1$ and thus $\psi(x^3 - f) = 1$. Symmetry implies $\psi(x^1 - f) = 1$.

iii) Without loss of generality, assume that $x = \mu'x^1 + (1 - \mu')x^3 = \mu''x^2 + (1 - \mu'')x^3$ for some $0 \leq \mu'' \leq 1$. Convexity of ψ then implies that $\psi(x - f) \leq \mu''\psi(x^2 - f) + (1 - \mu'')\psi(x^3 - f) = 1$. Applying ii) to the triplet x^1, x^2, x , we get $\psi(x - f) = 1$. \square

Lemma 2.3. *Let $f \in \mathbb{Q}^2$ and let ψ be valid for R_f . Assume that $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_j is valid and minimal for $j = 1, 2$.*

- i) *If $\psi(y - f) = 1$ for an integral point y then $\psi_j(y - f) = 1$ for $j = 1, 2$.*
- ii) *Let y^1 and y^2 two distinct integral points. Let E be a line segment with $y^1, y^2 \in E$ and with y^1 in the interior of E . If $\psi(x - f) = 1$ for all $x \in E$ then $\psi_j(x - f) = 1$ for all $x \in E$, $j = 1, 2$.*
- iii) *Let x^1 and y^1 be two distinct points, with y^1 integral. Let E be a line segment with $x^1, y^1 \in E$ and with y^1 in the interior of E . If $\psi(x - f) = 1$ for all $x \in E$ and $\psi_1(x^1 - f) = \psi_2(x^1 - f) = 1$ then $\psi_j(x - f) = 1$ for all $x \in E$, $j = 1, 2$.*
- iv) *Let L be a line with direction $d \in \mathbb{Q}^2$ and containing integral points. If $\psi(x - f) = 1$ for all $x \in L$, then $\psi_j(x - f) = 1$ for all $x \in L$, $j = 1, 2$.*

Proof. i) Since y is an integral point and ψ_j is valid, $\psi_j(y - f) \geq 1$. As $\psi(y - f) = 1$ and $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$, we get $\psi_j(y - f) = 1$ for $j = 1, 2$.

ii) Let $x \in E$ with x outside the line segment y^1y^2 with y^1 between x and y^2 . Such an x exists as y^1 is in the interior of E . We claim that $\psi_j(x - f) = 1$ for $j = 1, 2$. If x is integral, point i) proves it. Assume now that x is not integral. Lemma 2.2 i) for ψ_j and the triplet y^2, y^1, x shows that $\psi_j(x - f) \geq 1$. As $\psi(x - f) = 1$ and $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$, we get $\psi_j(x - f) = 1$. This proves the claim.

Using Lemma 2.2 iii), we get that $\psi_j(z - f) = 1$ for all z on the line segment xy^2 for $j = 1, 2$.

If there exists a point x outside the segment y^1y^2 with y^2 between x and y^1 , a similar reasoning can be used to get $\psi_j(z - f) = 1$ for all z on the line segment xy^1 for $j = 1, 2$.

iii) Similar to ii) with x^1 replacing y^2 .

iv) Let $\bar{x} \in L$. Observe that since $d \in \mathbb{Q}^2$, there exist infinitely many integral points on L . It is thus possible to find three distinct integral points $y^1, y^2, y^3 \in L$ such that both y^2 and \bar{x} are in the interior of the line segment $E = y^1y^3$. Since $\psi(x - f) = 1$ for all $x \in E$, point ii) above implies that $\psi_j(x - f) = 1$ for all $x \in E$ (and thus for \bar{x}), for $j = 1, 2$. \square

3 Facets of $R_f(r^1, \dots, r^k)$

In this section we prove Theorem 1.7. We first show that degenerate cases can be ignored when dealing with $R_f(r^1, \dots, r^k)$. We then characterize when triangle and quadrilateral inequalities define facets. Finally we give conditions for split inequalities to define facets of $R_f(r^1, \dots, r^k)$.

3.1 Degenerate cases are not needed for $R_f(r^1, \dots, r^k)$

Theorem 3.1. *Let $f, r^1, \dots, r^k \in \mathbb{Q}^2$ with $k \geq 1$. Every nontrivial facet of $R_f(r^1, \dots, r^k)$ can be obtained from a nondegenerate minimal valid function ψ .*

Proof. It suffices to show that any degenerate minimal valid function ψ is identical to a nondegenerate minimal valid function ψ' in the k directions r^1, \dots, r^k . Without loss of generality, assume that $\psi(r^i) < +\infty$ if and only if $i \leq \ell$ for some $0 \leq \ell \leq k$. We consider the five possible degenerate cases for ψ .

i) ψ defines a degenerate split inequality. Let L_0 be the infinite edge of $\text{cl}B_\psi$ containing f and let y^1 and y^2 be the first integral points on L_0 encountered starting from f in each direction, respectively. Let L_1 be the other infinite edge of $\text{cl}B_\psi$ and let y^3 and y^4 be two distinct integral points on L_1 . For $i = 1, \dots, \ell$, if there exists $\mu_i > 0$ such that $f + \mu_i r^i$ is a point on L_1 , let w^i be this point. Otherwise, define $w^i = f$. The convex hull of $\{w^i \mid i = 1, \dots, \ell\} \cup \{y^1, \dots, y^4\}$ is a lattice-free quadrilateral Q with no integral point in the interior of the edge E containing f . Replacing E by an edge parallel to E arbitrarily close to E , enlarging slightly Q gives a lattice-free quadrilateral Q' with f in its interior. Let ψ' be the minimal valid function corresponding to Q' . We have $\psi'(r^i) \leq \psi(r^i)$ for $i = 1, \dots, k$, proving the result.

ii) ψ defines a vertex-degenerate triangle or quadrilateral inequality. Let E_1 and E_2 be the two edges of $\text{cl}B_\psi$ incident with f and let y^1 and y^2 be the first integral points encountered on these edges starting from f . Let x be a point arbitrarily close to f with $\psi(x - f) = +\infty$ and such that the cone K with vertex x and generated by the vectors xy^1 and xy^2 contains all positive multiples of the rays r^1, \dots, r^ℓ except, possibly, those in directions fy^1 or fy^2 . If ψ defines a triangle inequality, let L_1 be the line supported by the edge of $\text{cl}B_\psi$ not containing f and let H be the half-space limited by L_1 and containing x . Otherwise, ψ defines a quadrilateral inequality, and let H be the cone formed by the two edges of the quadrilateral not containing f . The intersection of K with H is a lattice-free triangle T or quadrilateral Q with f in its interior. Let ψ' be the minimal function corresponding to T or Q . We have $\psi'(r^i) \leq \psi(r^i)$ for $i = 1, \dots, k$, proving the result.

iii) ψ defines an edge-degenerate triangle or quadrilateral inequality. Consider first the case where the edge E_1 containing f also contains an integral point interior to E_1 on each side

of f . This situation can only occur when $\text{cl}B_\psi$ is a triangle by Theorem 1.2 iv). Let y^1 and y^2 be the first integral points on E_1 encountered starting from f in each direction, respectively. Let x be a point arbitrarily close to f with $\psi(x - f) = +\infty$ and such that the cone K with vertex x and generated by the vectors xy^1 and xy^2 contains all the positive multiples of the rays r^1, \dots, r^ℓ except, possibly, those in directions fy^1 or fy^2 . The intersection of K with the cone formed by the two edges of $\text{cl}B_\psi$ not containing f is a lattice-free quadrilateral Q with f in the interior of Q . Let ψ' be the minimal valid function corresponding to Q . We have $\psi'(r^i) \leq \psi(r^i)$ for $i = 1, \dots, k$, proving the result.

Finally consider the case where the interior of the edge E_1 that contains f has all its integral points on one side of f . Let y be the closest integral point of f in the interior of E_1 . By tilting the edge E_1 around y by a small angle we modify the original triangle T (resp., quadrilateral Q) into a new triangle T' (resp. quadrilateral Q') with f in its interior. Let L'_1 be the line containing this tilted edge. Choose the tilting angle small enough so that all rays r^1, \dots, r^ℓ except, possibly, those in the direction fy point in the half-space limited by L'_1 and containing T' (resp., Q'). Let ψ' be the minimal valid function corresponding to T' (resp., Q'). We have $\psi'(r^i) \leq \psi(r^i)$ for $i = 1, \dots, k$, proving the result. \square

3.2 Triangle and quadrilateral inequalities for $R_f(r^1, \dots, r^k)$

Let us turn now to the nondegenerate cases, i.e. f is in the interior of B_ψ . We distinguish two cases depending on whether B_ψ is unbounded or a polytope. We deal with the polytopes in this section and with the unbounded case in the next section.

3.2.1 Reducing the dimension k

Let $f, r^1, \dots, r^k \in \mathbb{Q}^2$ with $k \geq 1$. Let B_ψ be a maximal lattice-free triangle or quadrilateral with f in its interior. In this section we reduce the question of whether ψ defines a facet of $R_f(r^1, \dots, r^k)$ to a problem on at most four rays r^j .

For $j = 1, \dots, k$, let p^j be the intersection of the half-line $f + \lambda r^j$, $\lambda \geq 0$, with the boundary of B_ψ . The point p^j is called the *boundary point* for r^j . Let P be a set of boundary points. We say that a point $p \in P$ is *active* if it can have a positive coefficient in a convex combination of points in P generating an integral point. Note that $p \in P$ is active if and only if p is integral or there exists $q \in P$ such that the segment pq contains an integral point in its interior. We say that an active point $p \in P$ is *uniquely active* if it has a positive coefficient in *exactly one* convex combination of points in P generating an integral point.

We apply the following *Reduction Algorithm*:

- 0.) Let $P = \{p^1, \dots, p^k\}$.
- 1.) While there exists $p \in P$ such that p is active and p is a convex combination of other points in P , remove p from P . At the end of this step, P contains at most two active points on each edge of B_ψ and all points of P are distinct.
- 2.) While there exists a uniquely active $p \in P$, remove p from P .
- 3.) If P contains exactly two active points p and q (and possibly inactive points), remove both p and q from P .

We say that *the ray condition holds* for a triangle or a quadrilateral if $P = \emptyset$ at termination of the Reduction Algorithm.

Lemma 3.2. *At termination of the Reduction Algorithm, the set of active points in P is either empty or consists of the corner points of B_ψ .*

Proof. Let Q be the set of active points in P at termination of the Reduction Algorithm. Suppose the lemma does not hold.

Observation 1: Step 1 implies that Q has at most two points on each edge.

Observation 2: Step 2 implies that every point of Q is involved in at least two distinct convex combinations.

Observation 3: Let E be an edge of B_ψ with a unique integral interior point. By Observations 1 and 2, Q cannot contain a point interior to E .

Observation 4: Step 3 and Observation 2 imply that $|Q| \geq 3$.

Suppose B_ψ is a triangle of Type i) or ii) of Lemma 2.1. Observation 3 and the fact that Q misses at least one corner of B_ψ implies that all the points of Q are on the same edge. This contradicts Observations 1 and 4.

Suppose B_ψ is a triangle of Type iii) of Lemma 2.1. By Observations 3 and 4, Q has two points in the edge with multiple integral points and one in the corner of B_ψ opposite that edge. This last point contradicts Observation 2 since Q misses at least one corner of B_ψ .

Suppose B_ψ is a quadrilateral. By Observation 3, the points of Q are corner points of B_ψ . Since Q misses at least one corner of B_ψ , there is a point $q \in Q$ such that only one of the two edges containing q also contains another point of Q . But then q contradicts Observation 2. \square

Let x^1, \dots, x^h denote the corner points of B_ψ . We define the *corner rays* of B_ψ to be the rays $r^j = x^j - f$ for $j = 1, \dots, h$.

Theorem 3.3. *Let $f, r^1, \dots, r^k \in \mathbb{Q}^2$ with $k \geq 1$. Let B_ψ be a maximal lattice-free triangle or quadrilateral with f in its interior. Let $h = 3$ when B_ψ is a triangle and $h = 4$ when B_ψ is a quadrilateral. Then ψ defines a facet of $R_f(r^1, \dots, r^k)$ if and only if either*

i) *the set $\{r^1, \dots, r^k\}$ contains rays r^{i_1}, \dots, r^{i_h} that are positive multiples of the corner rays of B_ψ and ψ defines a facet of $R_f(r^{i_1}, \dots, r^{i_h})$, or*

ii) *the ray condition holds.*

Proof. The function ψ defines a face of $R_f(r^1, \dots, r^k)$. We study its dimension. Let $S = \{s^1, \dots, s^t\}$ in $R_f(r^1, \dots, r^k)$ be a maximum cardinality set of affinely independent points that satisfy the inequality $\sum_{j=1}^k \psi(r^j) s_j \geq 1$ as an equality.

Let R be the $2 \times k$ matrix whose column j is r^j for $j = 1, \dots, k$. Let S be the $k \times t$ matrix whose column i is s^i for $i = 1, \dots, t$. Let D be the $k \times k$ diagonal matrix whose (j, j) entry is $\psi(r^j)$ for $j = 1, \dots, k$. Observe that D is invertible and let

$$\bar{R} = R \cdot D^{-1} \quad \text{and} \quad \bar{S} = D \cdot S .$$

Let us denote column j of \bar{R} by \bar{r}^j and column i of \bar{S} by \bar{s}^i . Observe that for $j = 1, \dots, k$ we have $\psi(\bar{r}^j) = 1$ implying that the point $p^j = f + \bar{r}^j$ is on the boundary of B_ψ . As

multiplying each component of all the vectors in S by a positive number does not change its affine dimension, the column set of \bar{S} has the same affine dimension as the column set of S .

Claim 1: For $i = 1, \dots, t$, the coordinates of the vector \bar{s}^i are the coefficients of a convex combination of the points in P that yields an integral point on the boundary of B_ψ .

Indeed, we have

$$\sum_{j=1}^k \bar{s}_j^i = \sum_{j=1}^k \psi(r^j) s_j^i = 1$$

Moreover, $\bar{s}^i \geq 0$ and $f + \sum_{j=1}^k r^j s_j^i \in \mathbb{Z}^2$. Thus

$$\sum_{j=1}^k p^j \bar{s}_j^i = \sum_{j=1}^k (f + r^j) \bar{s}_j^i = f + \sum_{j=1}^k r^j s_j^i \in \mathbb{Z}^2 .$$

Therefore the coordinates of \bar{s}^i are the coefficients of a convex combination of points in P that yields an integral point. Since the only integral points in B_ψ are on its boundary, the claim follows.

Claim 1 implies that the columns of \bar{S} are affinely independent if and only if they are linearly independent.

Claim 2: Suppose that the Reduction Algorithm removes point $p^k \in P$ in Step 1. If $k = 1$, then ψ defines a facet of $R_f(r^1, \dots, r^k)$. Otherwise, ψ defines a face of dimension w of $R_f(r^1, \dots, r^k)$ if and only if ψ defines a face of dimension $w - 1$ of $R_f(r^1, \dots, r^{k-1})$.

The boundary point p^k is a convex combination of other points in P , say $p^k = \sum_{j=1}^{k-1} \alpha_j p^j$. Define the k -vector v by

$$v_j = \begin{cases} \alpha_j & \text{for } j = 1, \dots, k-1 \\ -1 & \text{for } j = k \end{cases} .$$

As p^k is active, there exists a nonzero entry in row k of \bar{S} . We can assume without loss of generality that $\bar{s}_k^t > 0$. The vector $\bar{s}^{t+1} = \bar{s}^t + \bar{s}_k^t v$ is in $R_f(r^1, \dots, r^k)$ and satisfies the inequality $\sum_{j=1}^k \psi(r^j) s_j \geq 1$ as an equality. Therefore it belongs to the affine space generated by the columns of \bar{S} . The vectors \bar{s}^t, \bar{s}^{t+1} are linearly independent. It is thus possible to find a set \bar{S}' of t linearly independent vectors in $\{\bar{s}^1, \dots, \bar{s}^{t+1}\}$ and containing \bar{s}^t, \bar{s}^{t+1} . Then replacing \bar{s}^i by $\bar{s}^i + \bar{s}_k^i v$ for all $\bar{s}^i \in \bar{S}'$ except \bar{s}^t is just doing elementary column operations and does not change the rank, since $v = \frac{1}{\bar{s}_k^t} (\bar{s}^{t+1} - \bar{s}^t)$. As the resulting vectors have a zero entry in component k except $\bar{s}_k^t > 0$, by removing \bar{s}^t from \bar{S}' and deleting row k in \bar{S}' , we reduce by one both the dimension of the whole space and the dimension of the affine space spanned by \bar{S} . The converse is proved analogously, adding one zero component to the columns in \bar{S} and one new column in \bar{S} corresponding to a convex combination where p^k is active. The dimension of the space is increased by one as is the number of affinely independent points on the face. This proves the claim.

Claim 3: Suppose that the Reduction Algorithm removes point $p^k \in P$ in Step 2. If $k = 1$, then ψ defines a facet of $R_f(r^1, \dots, r^k)$. Otherwise, ψ defines a face of dimension w of $R_f(r^1, \dots, r^k)$ if and only if ψ defines a face of dimension $w - 1$ of $R_f(r^1, \dots, r^{k-1})$.

Point p^k is involved in a single convex combination \bar{s}^i . If $k = 1$, p^k is integral and the result clearly holds. Otherwise, removing \bar{s}^i and component k in \bar{S} , we reduce by one both the dimension of the whole space and the dimension of the affine space spanned by \bar{S} . The converse is proved as above. This proves the claim.

Claim 4: Suppose that $k = 2$ and the Reduction Algorithm removes points $p^1, p^2 \in P$ in Step 3. Then ψ defines a facet of $R_f(r^1, r^2)$.

Points p^1, p^2 are involved in two distinct convex combinations \bar{s}^1, \bar{s}^2 , and the result clearly holds.

Using Claims 2, 3 and 4 recursively, we get that ψ defines a facet of $R_f(r^1, \dots, r^k)$ if and only the ray condition holds or ψ defines a facet of $R_f(r^{i_1}, \dots, r^{i_{k'}})$ where $P' = \{p^{i_1}, \dots, p^{i_{k'}}\}$ is the set of remaining boundary points at termination of the Reduction Algorithm.

Let S' be the $k' \times t'$ matrix obtained from \bar{S} while applying the Reduction Algorithm. If some $p \in P'$ is not active, then the row of S' corresponding to p has only zero entries. This implies $t' < k'$, proving that ψ does not define a facet of $R_f(r^{i_1}, \dots, r^{i_{k'}})$. Therefore ψ defines a facet of $R_f(r^1, \dots, r^k)$ if and only if all points in P' are active and either the ray condition holds or ψ defines a facet of $R_f(r^{i_1}, \dots, r^{i_{k'}})$. Now the theorem follows from Lemma 3.2. \square

3.2.2 Rationality of the triangles and quadrilaterals that define facets of $R_f(r^1, \dots, r^k)$

We say that a triangle or quadrilateral is *rational* if its boundary lines have rational equations, or equivalently, if the coordinates of its corner points are rational.

Lemma 3.4. *Let $f, r^1, \dots, r^k \in \mathbb{Q}^2$. The maximal lattice-free triangles and quadrilaterals of Theorems 3.3 i) are rational.*

Proof. By a theorem of Meyer [16], the rationality of f, r^1, \dots, r^k implies that $R_f(r^1, \dots, r^k)$ is a rational polyhedron. Therefore its facets $\sum_{j=1}^k \psi(r^j)s_j \geq 1$ have rational coefficients $\psi(r^j)$. For triangles and quadrilateral inequalities, these coefficients are strictly positive. Therefore $x^j = f + \frac{1}{\psi(r^j)}r^j$ has rational coordinates for $j = 1, \dots, k$. In particular, the coordinates of the corner points are rational. \square

As usual, let $f, r^1, \dots, r^k \in \mathbb{Q}^2$. Suppose that B_ψ is a maximal lattice-free triangle or quadrilateral with f in its interior such that the ray condition holds. By Theorem 3.3, ψ defines a facet of $R_f(r^1, \dots, r^k)$. It may happen that B_ψ is irrational, but there exists a maximal lattice-free *rational* triangle or quadrilateral $B_{\psi'}$ defining the same facet of $R_f(r^1, \dots, r^k)$ as B_ψ . The proof of this claim is straightforward, since the only edges of B_ψ that can possibly be irrational have only one integral point y in their interior and no ray intersects them except possibly at y . Pivoting the edge slightly around y , we can make the edge rational.

3.2.3 Corner rays

In this section, we assume that B_ψ is a maximal lattice-free triangle or quadrilateral and that f is in its interior.

Let x^1, \dots, x^h be the vertices of B_ψ and let r^1, \dots, r^h be the corner rays of B_ψ with $r^j = x^j - f$. We always assume that the vertices are topologically ordered so that the edges of the boundary of B_ψ are convex combinations of x^i and x^{i+1} with indices taken modulo h . Let y^i be an integral point that can be obtained as a nontrivial convex combination of x^i and x^{i+1} for $i = 1, \dots, h$ (indices are always implicitly taken modulo h).

Define Y as the $2 \times h$ matrix whose column i is the vector y^i for $i = 1, \dots, h$ (Recall that $h = 3$ or 4). Define X as the $2 \times h$ matrix whose column i is the vector x^i for $i = 1, \dots, h$. Let S be the $h \times h$ matrix whose column i is the vector corresponding to the coefficients in the convex combination of x^i and x^{i+1} giving y^i for $i = 1, \dots, h$.

We then have

$$Y = X \cdot S \tag{5}$$

with

$$S = \begin{pmatrix} \alpha & 0 & 1 - \gamma \\ 1 - \alpha & \beta & 0 \\ 0 & 1 - \beta & \gamma \end{pmatrix} \quad \text{or} \quad S = \begin{pmatrix} \alpha & 0 & 0 & 1 - \delta \\ 1 - \alpha & \beta & 0 & 0 \\ 0 & 1 - \beta & \gamma & 0 \\ 0 & 0 & 1 - \gamma & \delta \end{pmatrix}$$

where α, β, γ and δ are all strictly between 0 and 1.

Since we are interested in the dimension of faces of polyhedra, which requires checking affine independence of points, we add a third row full of 1s to the matrices Y (resp. X) to obtain matrix \bar{Y} (resp., \bar{X}). Due to the specific form of the matrix S , we still have

$$\bar{Y} = \bar{X} \cdot S . \tag{6}$$

Let A be an $m \times n$ matrix. The *nullspace* of A is $\mathcal{N}(A) = \{x \in \mathbb{R}^n \mid Ax = 0\}$ and the *columnspace* of A is $\mathcal{C}(A) = \{z \in \mathbb{R}^m \mid z = Ax \text{ for some } x \in \mathbb{R}^n\}$.

The following three results are classical results of linear algebra [15]:

Lemma 3.5. *Let A be an $m \times n$ matrix and B be an $n \times p$ matrix. Then*

$$\text{rank}(A \cdot B) = \text{rank}(B) - \dim(\mathcal{N}(A) \cap \mathcal{C}(B)) .$$

Corollary 3.6. *Let A be an $m \times n$ matrix and B an $n \times p$ matrix. If $\text{rank}(A) = n$, then*

$$\text{rank}(A \cdot B) = \text{rank}(B) .$$

Proof. If $\text{rank}(A) = n$, then $\mathcal{N}(A) = \{0\}$ and has dimension 0. Applying Lemma 3.5 yields the result. \square

Corollary 3.7. *Let A be an $m \times n$ matrix and B an $n \times p$ matrix. Then*

$$\text{rank}(A \cdot B) \leq \min\{\text{rank}(A), \text{rank}(B)\} .$$

Proof. Apply Lemma 3.5 to $A \cdot B$ and its transpose. \square

Triangle inequalities

Theorem 3.8. *Triangle inequalities define facets of $R_f(r^1, r^2, r^3)$ when r^1, r^2, r^3 are the corner rays.*

Proof. Since $h = 3$ and B_ψ is a triangle, both \bar{Y} and \bar{X} have rank 3. By Corollary 3.6, S has rank 3 too. It implies that the columns of S are affinely independent. Since they all satisfy with equality the inequality $\sum_{i=1}^3 \psi(r^i) s_i \geq 1$, this inequality defines a facet of $R_f(r^1, r^2, r^3)$. \square

Putting together Theorems 3.3 and 3.8, we get:

Theorem 3.9. *A triangle inequality defines a facet of $R_f(r^1, \dots, r^k)$ if and only if one of the following holds:*

- i) *the set $\{r^1, \dots, r^k\}$ contains rays r^{i_1}, \dots, r^{i_3} that are positive multiples of the corner rays of the triangle;*
- ii) *the ray condition holds.*

Quadrilateral inequalities

When $k = 4$, both \bar{Y} and \bar{X} have rank 3. By Lemma 3.5, we have

$$3 = \text{rank}(\bar{Y}) = \text{rank}(\bar{X} \cdot S) = \text{rank}(S) - \dim(\mathcal{N}(\bar{X}) \cap \mathcal{C}(S)) .$$

Since $\text{rank}(\bar{X}) = 3$, we have that $\mathcal{N}(\bar{X})$ is a one-dimensional linear space. Hence $\dim(\mathcal{N}(\bar{X}) \cap \mathcal{C}(S)) \leq 1$ and $\text{rank}(S) = 4$ if and only if $\mathcal{N}(\bar{X}) \subseteq \mathcal{C}(S)$.

Theorem 3.10. *Consider a maximal lattice-free quadrilateral with vertices x^i , integral point y^i on edge $x^i x^{i+1}$ (indices taken modulo 4) and corner rays r^i , $i = 1, \dots, 4$. The corresponding quadrilateral inequality defines a facet of $R_f(r^1, r^2, r^3, r^4)$ if and only if there is no $t \in \mathbb{R}_+$ such that the point y^i divides the edge joining x^i to x^{i+1} in a ratio t for odd i and in a ratio $1/t$ for even i , i.e.*

$$\frac{\|y^i - x^i\|}{\|y^i - x^{i+1}\|} = \begin{cases} t & \text{for } i = 1, 3 \\ \frac{1}{t} & \text{for } i = 2, 4 \end{cases} . \quad (7)$$

Proof. Let F be the face of $R_f(r^1, \dots, r^4)$ defined by $\sum_{i=1}^4 \psi(r^i) s_i = 1$. As $f + r^i = x^i$ is on the boundary of B_ψ , we have $\psi(r^i) = 1$ for $i = 1, \dots, 4$. Hence, if $s \in F$ then $\sum_{i=1}^4 s_i = 1$. Recall that $R_f(r^1, \dots, r^k)$ is the convex hull of vectors in the set $H := \{s \in \mathbb{R}_+^4 \mid f + \sum_{i=1}^4 r^i s_i \text{ is integral}\}$. Thus, if F is a facet, then there exist four affinely independent vectors s^j , for $j = 1, \dots, 4$, in H with

$$\sum_{i=1}^4 s_i^j = 1 \quad \text{and} \quad z^j = f + \sum_{i=1}^4 r_i s_i^j = \sum_{i=1}^4 (f + r_i) s_i^j \quad \text{integral} .$$

This implies that z^j is in the convex hull of x^1, \dots, x^4 , for $j = 1, \dots, 4$. Theorem 1.2 shows that the only integral points in B_ψ are the points y^1, \dots, y^4 . Moreover, for each $j = 1, \dots, 4$, there is a unique convex combination of x^1, \dots, x^4 that produces y^j , namely column j of matrix S . In other words, F is a facet if and only if the columns of S are affinely independent. Observe that the columns of S are affinely independent if and only if they are linearly independent since the sum of the entries in any column of S is 1. It follows that F is a facet if and only if $\text{rank}(S) = 4$.

Let $u = (1, -1, 1, -1)^T$. By Theorem 1.2 iv), the points y^1, \dots, y^4 are the vertices of a parallelogram. This implies that $\bar{Y} \cdot u = 0$. Then (6) gives $\bar{X} \cdot S \cdot u = 0$. We now have two cases:

i) $S \cdot u = 0$. Then $\text{rank}(S) \leq 3$ and Corollary 3.7 shows that $\text{rank}(S) = 3$. Solving the linear system $S \cdot u = 0$ gives $\alpha = 1 - \beta = \gamma = 1 - \delta$. This is equivalent to the ratio condition of the statement.

ii) $S \cdot u \neq 0$. Then for $v = S \cdot u$ we have $\bar{X} \cdot v = 0$, and as $v \neq 0$, we have that $\mathcal{N}(\bar{X})$ is the linear space spanned by v . Since v is obtained as a linear combination of the columns of S , we have $\mathcal{N}(\bar{X}) \subseteq \mathcal{C}(S)$ and by Lemma 3.5 we get $\text{rank}(S) = 4$. Since all the columns of S satisfy with equality the inequality $\sum_{i=1}^4 \psi(r^i) s_i \geq 1$, this inequality defines a facet of $R_f(r^1, r^2, r^3, r^4)$. \square

We say that a maximal lattice-free quadrilateral *satisfies the ratio condition* when (7) does not hold for any $t > 0$.

We illustrate the ratio condition by a couple of examples. The quadrilateral inequality generated from the square whose edges contain the integral points $\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ in their middle does not define a facet of R_f , independently of the position of f in its interior. As mentioned in [4], when f is in the center of the square, the resulting quadrilateral inequality is a convex combination of the two split inequalities whose respective unbounded directions are the two coordinate axes. However, if one tilts just one edge of the square around its (integral) middle point, the resulting trapezoid has three distinct ratios $\frac{\|y^i - x^i\|}{\|y^i - x^{i+1}\|}$. Therefore the ratio condition is satisfied and Theorem 3.10 states that the resulting quadrilateral inequality defines a facet of R_f .

We give another more complicated example, see Figure 6. Let $f = \begin{pmatrix} \frac{1}{2} \\ \frac{1}{2} \end{pmatrix}$ and Q the quadrilateral with vertices $x^1 = \begin{pmatrix} \frac{7}{6} \\ \frac{1}{6} \end{pmatrix}, x^2 = \begin{pmatrix} \frac{7}{8} \\ \frac{13}{8} \end{pmatrix}, x^3 = \begin{pmatrix} -\frac{7}{6} \\ \frac{1}{6} \end{pmatrix}, x^4 = \begin{pmatrix} \frac{7}{8} \\ -\frac{1}{8} \end{pmatrix}$.

Edge x^1x^2 contains integral point $y^1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ with ratio $\frac{\|y^1 - x^1\|}{\|y^1 - x^2\|} = \frac{4}{3}$.

Edge x^2x^3 contains integral point $y^2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ with ratio $\frac{\|y^2 - x^2\|}{\|y^2 - x^3\|} = \frac{3}{4}$.

Edge x^3x^4 contains integral point $y^3 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ with ratio $\frac{\|y^3 - x^3\|}{\|y^3 - x^4\|} = \frac{4}{3}$.

Edge x^4x^1 contains integral point $y^4 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ with ratio $\frac{\|y^4 - x^4\|}{\|y^4 - x^1\|} = \frac{3}{4}$.

Theorem 3.10 states that the quadrilateral inequality obtained from Q is not a facet.




Figure 6: Illustration for the second example.

Indeed, it can be obtained as a convex combination of two triangle inequalities, each with a multiplier $\frac{1}{2}$. The first triangle T_1 has vertices $\begin{pmatrix} \frac{3}{2} \\ 0 \end{pmatrix}$, $\begin{pmatrix} \frac{4}{5} \\ \frac{7}{5} \end{pmatrix}$, $\begin{pmatrix} -2 \\ 0 \end{pmatrix}$. The second triangle has vertices $\begin{pmatrix} -\frac{1}{3} \\ -\frac{1}{3} \end{pmatrix}$, $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$, $\begin{pmatrix} -\frac{3}{4} \\ \frac{1}{4} \end{pmatrix}$. Both triangles have all four points y^1, y^2, y^3, y^4 on their boundaries. The corner rays of Q are $r^1 = \begin{pmatrix} \frac{2}{3} \\ -\frac{1}{3} \end{pmatrix}$, $r^2 = \begin{pmatrix} \frac{3}{8} \\ \frac{9}{8} \end{pmatrix}$, $r^3 = \begin{pmatrix} -\frac{5}{3} \\ -\frac{1}{3} \end{pmatrix}$, $r^4 = \begin{pmatrix} \frac{3}{8} \\ -\frac{5}{8} \end{pmatrix}$. Triangle T_1 has corner rays positive multiples of r^1, r^2 and r^3 . Triangle T_2 has corner rays positive multiples of r^2, r^3 and r^4 . If ψ, ψ_1 and ψ_2 denote the functions defined by Q, T_1 and T_2 respectively, it is easy to verify that $\psi = \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ in each of the cones $r^i r^{i+1}$ (indices defined modulo 4). Indeed, each of these functions is linear in each of the cones. So it is sufficient to verify the equality $\psi(r) = \frac{1}{2}\psi_1(r) + \frac{1}{2}\psi_2(r)$ in each of the directions $r^i, i = 1, \dots, 4$. In direction r^1 we have $\psi_1\left(\begin{pmatrix} 1 \\ -\frac{1}{2} \end{pmatrix}\right) = 1$ and $\psi_2\left(\begin{pmatrix} \frac{1}{2} \\ -\frac{1}{4} \end{pmatrix}\right) = 1$. This implies $\psi_1(r^1) = \frac{2}{3}$ and $\psi_2(r^1) = \frac{4}{3}$. Therefore $\psi(r^1) = \frac{1}{2}\psi_1(r^1) + \frac{1}{2}\psi_2(r^1)$ as required. Similarly, for the other rays, we find $\psi_1(r^2) = \frac{5}{4}$ and $\psi_2(r^2) = \frac{3}{4}$; $\psi_1(r^3) = \frac{4}{3}$ and $\psi_2(r^3) = \frac{2}{3}$; $\psi_1(r^4) = \frac{3}{4}$ and $\psi_2(r^4) = \frac{5}{4}$.

Putting together Theorems 3.3 and 3.10, we get:

Theorem 3.11. *A quadrilateral inequality defines a facet of $R_f(r^1, \dots, r^k)$ if and only if one of the following holds:*

- i) the set $\{r^1, \dots, r^k\}$ contains rays r^{i_1}, \dots, r^{i_4} that are positive multiples of the corner rays of the quadrilateral, and the ratio condition is satisfied;*
- ii) the ray condition holds.*

When the ray condition holds for a quadrilateral B_ψ , the half-lines $f + \lambda r^j$, $\lambda \geq 0$, can only intersect the boundary of B_ψ at integral points. When this happens, it is easy to find a maximal lattice-free triangle producing the same facet. Therefore we do not need to mention the ray condition in iii) of Theorem 1.7.

3.3 Split inequalities for $R_f(r^1, \dots, r^k)$

When B_ψ is unbounded, ψ defines a split inequality for R_f by Theorem 1.1. We will see in Theorem 4.1 that split inequalities always define facets of R_f . The situation for $R_f(r^1, \dots, r^k)$ is a little bit more complicated, as the next theorem shows.

When $\psi(r^i) > 0$ for $i = 1, \dots, k$, we can define the boundary point for r^i similarly to the case of triangles and quadrilaterals, and we can apply the Reduction Algorithm of Section 3.2.1. We say that *the ray condition holds* for a split if, at termination of the Reduction Algorithm, either $P = \emptyset$, or $P = \{p_1, q_1, p_2, q_2\}$ with p_1, q_1 on one of the boundary lines and p_2, q_2 on the other and both line segments p_1q_1 and p_2q_2 contain at least two integral points.

Theorem 3.12. *Let $f, r^1, \dots, r^k \in \mathbb{Q}^2$ with $k \geq 1$. Let ψ define a split inequality. The inequality $\sum_{i=1}^k \psi(r^i) s_i \geq 1$ defines a facet of $R_f(r^1, \dots, r^k)$ if and only if*

- i) $\psi(r^i) = 0$ for some $i = 1, \dots, k$; or
- ii) $\psi(r^i) > 0$ for all $i = 1, \dots, k$ and the ray condition holds.

Proof. Suppose that $\psi(r^i) = 0$ for some $i \in \{1, \dots, k\}$. By Theorem 1.1, the only minimal valid inequality having the coefficient of s_i equal to 0 is the split inequality defined by ψ . Therefore $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_1 and ψ_2 are both valid and minimal implies that $\psi = \psi_1 = \psi_2$. Remark 1.4 proves that ψ is a facet of $R_f(r^1, \dots, r^k)$.

Suppose now that $\psi(r^i) > 0$ for $i = 1, \dots, k$. Using arguments similar to those used in the proof of Lemma 3.2, we obtain that at termination of the Reduction Algorithm, the set Q of active points is either empty, or it contains exactly two points p_1, q_1 in L_1 and two points p_2, q_2 in L_2 where L_1, L_2 are the boundary lines of B_ψ , and both line segments p_1q_1 and p_2q_2 contain at least two integral points. Using arguments similar to those used in the proof of Theorem 3.3, we obtain that ψ defines a facet if and only if, at termination of the Reduction Algorithm, the set P is empty or only contains active points. In other words, ψ defines a facet if and only if the ray condition holds. \square

4 Facets of R_f

4.1 Split inequalities define facets of R_f

4.1.1 Nondegenerate case

Let $f \in \mathbb{Q}^2$. Consider a direction $r^0 \in \mathbb{Q}^2 \setminus \{0\}$ such that the line $L_0 := \{x = f + \alpha r^0, \alpha \in \mathbb{R}\}$ contains no integral point. Let L_1 and L_2 be parallel lines to L_0 , each containing integral points, such that the set of points between L_1 and L_2 contains no integral point in its interior and contains L_0 . (See Figure 7.) Define $\psi(r_0) = \psi(-r_0) = 0$, $\psi(x - f) = 1$ for any $x \in L_1 \cup L_2$.

Theorem 4.1. *Let $f \in \mathbb{Q}^2$. Split inequalities define facets of R_f .*




Figure 7: Illustration for Theorem 4.1.

Proof. Suppose not. By Remark 1.4, $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_1, ψ_2 are minimal valid functions and $\psi \neq \psi_1$ or ψ_2 .

By Theorem 1.1, $\psi_j(r_0) \geq 0$ for $j = 1, 2$. Since $\psi(r_0) = 0$, we have $\psi_j(r_0) = 0$ for $j = 1, 2$. Similarly $\psi_j(-r_0) = 0$.

Lemma 2.3 iv) for L_i proves that $\psi_j(x - f) = 1$ for all $x \in L_i$, for all $i = 1, 2, j = 1, 2$.

By homogeneity, ψ_j is therefore defined over all of \mathbb{R}^2 . But then $\psi_1 = \psi_2 = \psi$, a contradiction. \square

4.1.2 Degenerate case

Consider a direction $r^0 \in \mathbb{Q}^2 \setminus \{0\}$ such that the line $L_0 := \{x = f + \alpha r^0, \alpha \in \mathbb{R}\}$ contains integral points. Let L_1 be a line parallel to L_0 that contains integral points, such that the set of points between L_0 and L_1 contains no integral point in its interior. Let y^1 and y^2 be the first integral points encountered on the half-lines $f + \alpha r^0, \alpha \geq 0$, and $f - \alpha r^0, \alpha \geq 0$ respectively. (See Figure 8.) Define $\psi(y^1 - f) = \psi(y^2 - f) = 1$ and $\psi(x - f) = 1$ for any $x \in L_1$. Since ψ is homogeneous, this defines $\psi(r)$ for all $r \in \mathbb{Q}^2$ in the closed half-space limited by L_0 and containing L_1 . For all other $r \in \mathbb{Q}^2 \setminus \{0\}$, define $\psi(r) = +\infty$. The inequality $\sum \psi(r) s_r \geq 1$, a *degenerate split inequality*, is valid for R_f .




Figure 8: Illustration for Theorem 4.2.

Theorem 4.2. Let $f \in \mathbb{Q}^2$. Degenerate split inequalities define facets of R_f .

Proof. Suppose not. Then $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_1, ψ_2 are minimal valid functions and $\psi \neq \psi_1$ or ψ_2 .

Lemma 2.3 i) proves that $\psi_j(y^1 - f) = 1$ for $j = 1, 2$ and Lemma 2.3 iv) for L_1 proves that $\psi_j(x - f) = 1$ for all $x \in L_1, j = 1, 2$.

By homogeneity, $\psi_j(x - f)$ is therefore defined for all $x \in \mathbb{R}^2$ that are in the closed half-space H limited by L_0 and containing L_1 . Observe that $\psi(x - f) = \psi_j(x - f)$ for all $x \in \mathbb{R}^2$ and in H .

Suppose that there exists $x^1 \notin H$ with $\psi_j(x^1 - f) < +\infty$ for $j = 1$ or $j = 2$. Then $\psi_j(x - f) \leq \psi(x - f)$ for all $x \in \mathbb{R}^2$ with strict inequality for x^1 , a contradiction to the minimality of ψ .

It follows that $\psi_1 = \psi_2 = \psi$, a contradiction. \square

4.2 Nondegenerate triangle and quadrilateral inequalities

Let $f \in \mathbb{Q}^2$. Nondegenerate minimal valid inequalities that are not split inequalities are generated by a function ψ such that B_ψ is either a triangle or a quadrilateral and f is in the interior of B_ψ . Let x^1, \dots, x^k be the vertices of B_ψ with $k = 3$ or $k = 4$. Note that these vertices may have irrational coordinates. Let $\{r^1, \dots, r^k\}$ be the corner rays of B_ψ , namely $r^i = x^i - f$.

Given a valid and minimal function ψ for R_f and rays r^1, \dots, r^k we say that two valid and minimal functions ψ_1 and ψ_2 for R_f *dominate* ψ *restricted to* $\{r^1, \dots, r^k\}$ if $\psi(r^i) \geq \frac{1}{2}\psi_1(r^i) + \frac{1}{2}\psi_2(r^i)$ for $i = 1, \dots, k$ and $\psi(r^i) = \psi_1(r^i)$ and $\psi(r^i) = \psi_2(r^i)$ does not hold for all $i = 1, \dots, k$.

The next theorem will allow us to extend the proof of earlier results to cover the case of possibly irrational corner rays.

Theorem 4.3. *Let $f \in \mathbb{Q}^2$ and let ψ be a minimal valid function for R_f . Assume that B_ψ is a polytope with f in its interior and let r^1, \dots, r^k be the corner rays of B_ψ . Then there exist two minimal valid functions ψ_1 and ψ_2 dominating ψ restricted to $\{r^1, \dots, r^k\}$ if and only if ψ is not facet defining for R_f .*

Proof. Let x^1, \dots, x^k be the vertices of B_ψ . Let E_i denote the edge of B_ψ between vertices x^i and x^{i+1} for $i = 1, \dots, k$ and let y^i be an integral point that can be obtained as a nontrivial convex combination of x^i and x^{i+1} for $i = 1, \dots, k$ (indices are implicitly taken modulo k).

Assume first that the two functions ψ_1, ψ_2 of the statement of the theorem exist. Let $\bar{\psi} = \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$. Consider the edge joining x^i to x^{i+1} . Since ψ_1 and ψ_2 are valid functions and y^i is integral, we have $\psi_1(y^i - f) \geq 1$ and $\psi_2(y^i - f) \geq 1$. This implies $\bar{\psi}(y^i - f) \geq 1$. On the other hand, convexity of $\bar{\psi}$ implies $\bar{\psi}(y^i - f) \leq \lambda\bar{\psi}(x^i - f) + (1 - \lambda)\bar{\psi}(x^{i+1} - f)$ where $0 < \lambda < 1$ is the convex combination of x^i, x^{i+1} producing y^i . Since ψ_1 and ψ_2 dominate ψ restricted to $\{r^1, \dots, r^k\}$, we get $\bar{\psi}(y^i - f) \leq \lambda\psi(x^i - f) + (1 - \lambda)\psi(x^{i+1} - f) = 1$. Therefore $\bar{\psi}(y^i - f) = 1$. Applying Lemma 2.2 i) to $\bar{\psi}$ and the triple $x^i y^i x^{i+1}$, we get $\bar{\psi}(r^{i+1}) \geq 1$. But as $1 = \psi(r^{i+1}) \geq \bar{\psi}(r^{i+1})$, we get $\bar{\psi}(r^{i+1}) = 1$. Applying Lemma 2.2 iii) to ψ and the triple $x^i y^i x^{i+1}$, we get that $\bar{\psi}(x - f) = \psi(x - f)$ for all x in the segment $x^i x^{i+1}$. A similar reasoning shows that $\bar{\psi}(x - f) = 1$ for all x on the boundary of B_ψ , proving that $\bar{\psi} = \psi$. But then ψ does not define a facet of R_f as it is a convex combination of ψ_1 and ψ_2 .

For the converse, assume that ψ does not define a facet of R_f and let $\psi_1 \neq \psi_2$ be two valid and minimal functions such that $\psi \geq \frac{1}{2} \psi_1 + \frac{1}{2} \psi_2$ with $\psi_1(z - f) \neq 1$ for some $z \in B_\psi$. Then either $\psi_1(z - f) < 1$ or $\psi_2(z - f) < 1$. Assume without loss of generality that $\psi_1(z - f) < 1$ and that $z \in E_i$. Applying Lemma 2.3 i) to y^i , we get $\psi_1(y^i - f) = 1$. If we have $\psi_1(r^i) = \psi_1(r^{i+1}) = 1$, applying Lemma 2.2 iii) to ψ_1 and the triple $x^i y^i x^{i+1}$, we get that $\psi_1(x - f) = 1$ for all x in the segment $x^i x^{i+1}$. This is a contradiction with $\psi_1(z - f) < 1$. Hence, either $\psi_1(r^i) \neq \psi(r^i) = 1$ or $\psi_1(r^{i+1}) \neq \psi(r^{i+1}) = 1$ and ψ_1, ψ_2 are two valid minimal functions dominating ψ restricted to $\{r^1, \dots, r^k\}$. \square

The following is an extension of Theorem 3.8 to cover the case of triangles with possibly irrational corner rays.

Lemma 4.4. *Let ψ be valid and minimal such that B_ψ is a triangle with corner rays r^1, r^2, r^3 and f in its interior. Then ψ restricted to $\{r^1, r^2, r^3\}$ cannot be dominated by two valid and minimal functions ψ_1 and ψ_2 .*

Proof. The proof of Theorem 3.8 can be used as is. The definitions of \bar{Y} , \bar{X} and S are not affected by the possible irrationality of some of the rays. It shows that the rank of S is 3, i.e. the face F of $\{s \in \mathbb{R}^3 \mid f + \sum_{j=1}^3 r^j s_j \in \mathbb{Z}^2, s \geq 0\}$ defined by $\sum \psi(r^j) s_j \geq 1$ has dimension 2. Its equation is uniquely defined. This implies that ψ restricted to $\{r^1, r^2, r^3\}$ cannot be dominated by two valid and minimal functions ψ_1, ψ_2 . \square

Similarly, the following is an extension of Theorem 3.10 to cover the case of quadrilaterals with possibly irrational corner rays.

Lemma 4.5. *Let ψ be valid and minimal such that B_ψ is a quadrilateral with corner rays r^1, r^2, r^3, r^4 and f in its interior. Then ψ restricted to $\{r^1, r^2, r^3, r^4\}$ cannot be dominated by two valid and minimal functions ψ_1 and ψ_2 if and only if the quadrilateral satisfies the ratio condition.*

Proof. The proof of Theorem 3.10 can be used as is. The definitions of \bar{Y} , \bar{X} and S are not affected by the possible irrationality of some of the rays. It shows that, depending on the ratio condition being satisfied or not, the rank of S is 4 or 3, i.e. the face F of $\{s \in \mathbb{R}^4 \mid f + \sum_{j=1}^4 r^j s_j \in \mathbb{Z}^2, s \geq 0\}$ defined by $\sum \psi(r^j) s_j \geq 1$ has dimension 3 or 2.

If the ratio condition is satisfied, then a proof similar to the one for Lemma 4.4 shows that ψ restricted to $\{r^1, r^2, r^3, r^4\}$ cannot be dominated by two valid and minimal functions.

If the ratio condition is not satisfied, there exists an hyperplane containing $F \cup \{0\}$ with equation $\sum_{j=1}^4 h_j s_j = 0$. For $\epsilon > 0$, define ψ_1, ψ_2 by $\psi_1(r^j) = \psi(r^j) + \epsilon h_j$ and $\psi_2(r^j) = \psi(r^j) - \epsilon h_j$. As ψ is valid, we have $\psi(y - f) \geq 1$ for all $y \in \mathbb{Z}^2$ and Theorem 1.2 iv) implies that equality holds only for the four integral points on the boundary of B_ψ , that is the four integral points obtained according to the convex combinations corresponding to the columns of S . This shows that for $\epsilon > 0$ small enough, the two functions ψ_1, ψ_2 can be seen as restrictions of two valid quadrilateral inequalities. As $\psi(r^j) = \frac{1}{2} \psi_1(r^j) + \frac{1}{2} \psi_2(r^j)$ for $j = 1, \dots, 4$, it follows that ψ restricted to $\{r^1, r^2, r^3, r^4\}$ is dominated by two valid and minimal functions. \square

As a consequence of the last three lemmas, we immediately get the following two results.

Theorem 4.6. *All maximal lattice-free triangles with f in their interior define facets of R_f .*

Theorem 4.7. *A maximal lattice-free quadrilateral having f in its interior defines a facet of R_f if and only if it satisfies the ratio condition.*

Although only rational triangles and quadrilaterals are needed to define the facets of $R_f(r^1, \dots, r^k)$ when f and the rays r^j 's are in \mathbb{Q}^2 (see Section 3.2.2), irrational triangles and quadrilaterals are needed to define some of the facets of R_f for $f \in \mathbb{Q}^2$.

4.3 Degenerate triangle inequalities

In this section, we use the following notation:

Notation 4.8. *Let T be a maximal lattice-free triangle in \mathbb{R}^2 . Let x^1, x^2, x^3 be the vertices of T such that edge E_i of T is the convex combination of x^i and x^{i+1} for $i = 1, 2, 3$ (indices are taken modulo 3). Let y^i be an integral point in the interior of E_i for $i = 1, 2, 3$.*

Note that the existence of y^i for $i = 1, 2, 3$ follows from Theorem 1.2.

4.3.1 f is a vertex of the triangle

Here, we assume also that f is a vertex of T . Without loss of generality, we assume $f = x^2$. Then edges E_1 and E_2 of T contain f . (See Figure 9.)

Define $\psi(y^1 - f) = \psi(y^2 - f) = 1$. By homogeneity, this defines $\psi(x - f)$ for all x on the half-lines L_1 and L_2 starting from f and containing y^1 and y^2 respectively. Define $\psi(x - f) = 1$ for any x in $E_3 \setminus \{E_1 \cup E_2\}$. By homogeneity, this defines ψ in the open cone C limited by L_1 and L_2 . Finally, define $\psi(x - f) = +\infty$ for any x outside $C \cup L_1 \cup L_2$. Then ψ defines a *vertex-degenerate triangle inequality*.




Figure 9: Illustration for Theorem 4.9.

Theorem 4.9. *A vertex-degenerate triangle inequality defines a facet of R_f if and only if the edge of T opposite f contains at least two integral points in its interior.*

Proof. Suppose first that the edge E_3 contains only one integral point y^3 in its interior. Construct maximal lattice-free triangles T' and T'' from T by keeping the half-lines L_1 and L_2 unchanged, and tilting the edge E_3 around the integral point y^3 so that the new edges E'_1 (E''_1 respectively) are slightly shorter (longer respectively) than E_1 . The same construction used to define the function ψ from T can be used to define functions ψ' and ψ'' from T' and

T'' respectively. The functions ψ' and ψ'' are linear in the open cone C limited by L_1 and L_2 and $\psi(y^3) = \psi'(y^3) = \psi''(y^3) = 1$. Therefore, we can write ψ as a convex combination of ψ' and ψ'' . This shows that ψ does not define a facet in this case.

Now consider the case where E_3 contains two integral points y^3 and y^4 in its interior. Suppose that ψ does not define a facet of R_f . Then $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_j is valid and minimal for $j = 1, 2$. Lemma 2.3 ii) shows that $\psi_j(x - f) = 1$ for all x in the interior of E_3 .

By Lemma 2.3 i), $\psi_j(y^1 - f) = 1$, for $j = 1, 2$. By homogeneity, this defines $\psi_j(x - f)$ for all x on the half-line L_1 . Similarly $\psi_j(y^2 - f) = 1$ and by homogeneity, this defines $\psi_j(x - f)$ for all x on the half-line L_2 . Observe that $\psi(x - f) = \psi_j(x - f)$ for all $x \in \mathbb{R}^2$ and in $C \cup L_1 \cup L_2$.

Suppose that there exists $x^4 \notin C \cup L_1 \cup L_2$ with $\psi_j(x^4 - f) < +\infty$ for $j = 1$ or $j = 2$. Then $\psi_j(x - f) \leq \psi(x - f)$ for all $x \in \mathbb{R}^2$ with strict inequality for x^4 , a contradiction with the minimality of ψ . It follows that $\psi_1 = \psi_2 = \psi$, a contradiction. Therefore ψ define a facet of R_f . \square

4.3.2 f is on an edge of the triangle

Here we use Notation 4.8 and we assume that f is in the interior of an edge of T . Without loss of generality, we assume $f \in E_1$ and that y^1 is between f and x^1 . (See Figure 10.)

Define $\psi(x - f) = 1$ for any $x \in (E_2 \setminus \{x^2\}) \cup (E_3 \setminus \{x^1\})$. Let L_1 denote the line containing the segment E_1 . Since ψ is homogeneous, $\psi(x - f)$ is defined for all $x \in \mathbb{R}^2$ in the open half-space H limited by L_1 and containing x^3 .

Define $\psi(y^1 - f) = 1$. On the half-line from f that goes through x^2 , let y^4 be the first integral point encountered, starting from f . If y^4 is reached before x^2 , then set $\psi(y^4 - f) = 1$. Otherwise set $\psi(x^2 - f) = 1$. By homogeneity, this defines $\psi(x - f)$ for all $x \in L_1$.

For all $x \in \mathbb{R}^2 \setminus (L_1 \cup H)$, define $\psi(x - f) = +\infty$. Then ψ defines an *edge-degenerate triangle inequality*.




Figure 10: Illustration for Theorem 4.10.

Theorem 4.10. *An edge-degenerate triangle inequality defines a facet of R_f if and only if at least one of the two edges not containing f contains at least two integral points y with $\psi(y - f) = 1$.*

Proof. Suppose first that E_2 and E_3 each contain only one integral point, y^2 and y^3 respectively. Let x^4 and x^5 be points on the line fx^3 such that x^3 is their middle point and the

distance between x^4 and x^5 is very small. Let T' (resp. T'') denote the triangle with one side contained in L_1 , one side containing x^4y^2 (resp. x^5y^2) and one side containing x^4y^3 (resp. x^5y^3). The convex sets T' and T'' define valid functions ψ' and ψ'' as above. Furthermore ψ is a convex combination of ψ' and ψ'' , showing that ψ is not facet defining.

Suppose now that edge E_i contains at least two integral points for $i = 2$ or 3 . Without loss of generality, we can assume $i = 2$. Suppose that ψ does not define a facet of R_f . Then $\psi \geq \frac{1}{2}\psi_1 + \frac{1}{2}\psi_2$ where ψ_j are valid and minimal for $j = 1, 2$. Lemma 2.3 ii) shows that $\psi_j(x - f) = 1$ for all $x \in E_2 \setminus E_1$. In particular $\psi_j(x^3 - f) = 1$ for $j = 1, 2$.

The edge E_3 contains the integral point y^3 in its interior and the point x^3 . Lemma 2.3 iii) shows that $\psi_j(x - f) = 1$ for all $x \in E_3 \setminus E_1$. This shows that $\psi_1(x - f) = \psi_2(x - f) = \psi(x - f)$ for all $x \in \mathbb{R}^2$ in the open half-space H containing x^3 limited by L_1 .

Lemma 2.3 i) shows that $\psi_j(y^1 - f) = 1$, for $j = 1, 2$. On the half-line of L_1 going through x^2 there are two cases. If an integral point y^4 is encountered starting from f before reaching x^2 , Lemma 2.3 i) yields $\psi_j(y^4 - f) = 1$, for $j = 1, 2$. So suppose that no integral point is encountered starting from f before reaching x^2 . In this case $\psi(x^2) = 1$ and Lemma 2.3 iii) for E_2 and with x^3 playing the role of x^1 of the lemma shows that we have $\psi_j(x^2) = 1$. By homogeneity, this defines $\psi_j(x - f)$ for all $x \in L_1$, for $j = 1, 2$.

Finally, the discontinuity of ψ_j on the half-line of L_1 from f in the direction of y^1 implies that $\psi_j(x - f) = +\infty$ for all $x \notin H \cup L_1$, using a similar argument as in the proof of Theorem 4.9. \square

4.4 Degenerate quadrilateral inequalities

In this section, we use the following notation:

Notation 4.11. Let Q be a maximal lattice-free quadrilateral in \mathbb{R}^2 . Let x^1, \dots, x^4 be the vertices of Q such that edge E_i of Q is the convex combination of x^i and x^{i+1} for $i = 1, \dots, 4$ (indices are taken modulo 4). Let y^i be the unique integral point in the interior of E_i for $i = 1, \dots, 4$.

Note that the existence and unicity of y^i for $i = 1, \dots, 4$ follows from Theorem 1.2.

4.4.1 f is a vertex of the quadrilateral

Here, we assume that f is a vertex of Q . Without loss of generality, we assume $f = x^2$. (See Figure 11.)

Define $\psi(y^1 - f) = \psi(y^2 - f) = 1$. By homogeneity, this defines $\psi(x - f)$ for all x on the half-lines L_1 and L_2 starting from f and containing y^1 and y^2 respectively. Define $\psi(x - f) = 1$ for any x in $E_3 \cup E_4 \setminus \{x^1 \cup x^3\}$. By homogeneity, this defines ψ in the open cone C limited by L_1 and L_2 . Finally, define $\psi(x - f) = +\infty$ for any x outside $C \cup L_1 \cup L_2$. Then ψ defines a *vertex-degenerate quadrilateral inequality*.

Theorem 4.12. A *vertex-degenerate quadrilateral inequality* never defines a facet of R_f .

Proof. Let x^5 and x^6 be points on the line fx^4 such that x^4 is their middle point and the distance between x^5 and x^6 is very small. Let Q' denote the quadrilateral with vertices x^5 , f and two others on lines fx^1 and fx^2 respectively, and containing the integral points y^i for




Figure 11: Illustration for Theorem 4.12.

$i = 1, \dots, 4$ in the interior of its edges. Quadrilateral Q'' is defined similarly except that it has vertex x^6 instead of x^5 . The convex sets Q' and Q'' define valid functions ψ' and ψ'' as above. Furthermore ψ is a convex combination of ψ' and ψ'' with $\psi_1 \neq \psi_2$, proving that the vertex-degenerate inequality is not a facet. \square

4.4.2 f is on an edge of the quadrilateral

Here, we use Notation 4.11 and we assume that f is on one of the edges of Q . Without loss of generality, we assume $f \in E_1$ and that y^1 is between f and x^1 .

Define $\psi(y^1 - f) = 1$. By homogeneity, this defines $\psi(x - f)$ for all x on the half-line of L_1 starting from f and containing y^1 . Define $\psi(x - f) = 1$ for any x in $E_2 \cup E_3 \cup E_4 \setminus \{x^1\}$. By homogeneity, this defines ψ in the rest of the closed half-plane H limited by the line L_1 . Finally, define $\psi(x - f) = +\infty$ for any x outside C . Then ψ defines an *edge-degenerate quadrilateral inequality*.

Theorem 4.13. *An edge-degenerate quadrilateral inequality never defines a facet of R_f .*

Proof. As in the proof of Theorem 4.12, we can perturb Q into two quadrilaterals Q' and Q'' going through f and y^i for $i = 1, \dots, 4$ and vertices on the lines fx^3 or fx^4 and construct the corresponding functions ψ' and ψ'' as above. Then ψ is a convex combination of ψ' and ψ'' proving that the edge-degenerate inequality is not a facet. \square

Acknowledgements: We would like to thank Kent Andersen, Santanu Dey, Yanjun Li and Giacomo Zambelli for their helpful comments. In particular Kent Andersen's comments helped us bring into sharper focus the results for $R_f(r^1, \dots, r^k)$ and Santanu Dey brought to our attention the importance of irrational polytopes for R_f .

References

- [1] K. Andersen, Q. Louveaux, R. Weismantel and L. Wolsey, Cutting Planes from Two Rows of a Simplex Tableau, *Proceedings of IPCO XII*, Ithaca, New York (June 2007) 1-15.

- [2] E. Balas, Intersection Cuts - A New Type of Cutting Planes for Integer Programming, *Operations Research* 19 (1971) 19-39.
- [3] D.E. Bell, A Theorem Concerning the Integer Lattice, *Studies in Applied Mathematics* 56 (1977) 187-188.
- [4] V. Borozan and G. Cornuéjols, Minimal Valid Inequalities for Integer Constraints, technical report (July 2007).
- [5] G. Cornuéjols, Y. Li, On the Rank of Mixed 0,1 Polyhedra, *Mathematical Programming A* 91 (2002) 391-397.
- [6] W. Cook, R. Kannan and A. Schrijver, Chvátal Closures for Mixed Integer Programming Problems, *Mathematical Programming* 47 (1990) 155-174.
- [7] S.S. Dey and J.-P. P. Richard, L. A. Miller and Y. Li, Extreme Inequalities for Infinite Group Problems, technical report (2006).
- [8] S.S. Dey and L.A. Wolsey, Lifting Integer Variables in Minimal Inequalities Corresponding to Lattice-Free Triangles, *IPCO XIII*, Bertinoro, Italy (June 2008).
- [9] R.E. Gomory, Some Polyhedra Related to Combinatorial Problems, *Journal of Linear Algebra and its Applications* 2 (1969) 451-558.
- [10] R.E. Gomory, An Algorithm for Integer Solutions to Linear Programs, *Recent Advances in Mathematical Programming*, R.L. Graves and P. Wolfe eds., McGraw-Hill, New York (1963) 269-302.
- [11] R.E. Gomory, Thoughts about Integer Programming, 50th Anniversary Symposium of OR, University of Montreal, January 2007, and Corner Polyhedra and Two-Equation Cutting Planes, George Nemhauser Symposium, Atlanta, July 2007.
- [12] R.E. Gomory and E.L. Johnson, Some Continuous Functions Related to Corner Polyhedra, Part I, *Mathematical Programming* 3 (1972) 23-85.
- [13] E.L. Johnson, On the Group Problem for Mixed Integer Programming, *Mathematical Programming Study* (1974) 137-179.
- [14] L. Lovász, Geometry of Numbers and Integer Programming, *Mathematical Programming: Recent Developments and Applications*, M. Iri and K. Tanabe eds., Kluwer (1989) 177-210.
- [15] C.D. Meyer, *Matrix Analysis and Applied Linear Algebra*, SIAM (2000).
- [16] R.R. Meyer, On the Existence of Optimal Solutions to Integer and Mixed Integer Programming Problems, *Mathematical Programming* 7 (1974) 223-235.
- [17] G.L. Nemhauser and L.A. Wolsey, *Integer and Combinatorial Optimization*, Wiley (1988).
- [18] H.E. Scarf, An Observation on the Structure of Production Sets with Indivisibilities, *Proc. Nat. Acad. Sciences USA* 74 (1977) 3637-3641.